CLAIMS

We Claim:

10

15

20

The first time after the first time that the first time time time the first time time time time time time time

- 5 1. A method for selecting an optimal coupling temperature of a nucleic acid synthesis reaction comprising:
 - a) providing:
 - i. a nucleic acid synthesizer comprising a heating component; and
 - ii. nucleic acid synthesis reagents;
 - b) synthesizing a plurality of nucleic acid molecules with said synthesizer and said nucleic acid synthesis reagents, wherein the synthesis of each nucleic acid molecule of said plurality of nucleic acid molecules comprises a coupling reaction performed at a different temperature within a temperature range; and
 - c) measuring nucleic acid synthesis efficiency for each of said syntheses of said plurality of nucleic acid molecules; and
 - d) selecting an optimal coupling temperature within said temperature range.
 - 2. The method of Claim 1, wherein said temperature range is 20 to 60 degrees C.
- A nucleic acid synthesizer comprising one or more reaction chambers and
 a heating component configured to heat said one or more reaction chambers during a synthesis reaction.
 - 4. The nucleic acid synthesizer of Claim 3, wherein said heating component comprises a resistance heater.

30

15

- 5. The nucleic acid synthesizer of Claim 3, wherein said heating component comprises a Peltier device.
- 6. The nucleic acid synthesizer of Claim 3, wherein said heating component5 comprises a heated reagent.
 - 7. The nucleic acid synthesizer of Claim 3, wherein said heating component comprises a magnetic induction device.
- 10 8. The nucleic acid synthesizer of Claim 3, wherein said heating component comprises microwaves.
 - 9. The nucleic acid synthesizer of Claim 3, wherein said heating component comprises a transfer of heat from a fluid or a gas.
 - 10. A nucleic acid synthesizer, comprising:
 - a. one or more reaction chambers containing an oligonucleotide; and
 - b. a heating component,

wherein said heating component is configured to heat said one or more reaction chambers during a synthesis reaction wherein said oligonucleotide is coupled to a synthesis reagent.

25

20